CORRECTION

Correction: A compound downregulation of *SRRM2* and miR-27a-3p with upregulation of miR-27b-3p in PBMCs of Parkinson's patients is associated with the early stage onset of disease

Soudabeh Fazeli, Majid Motovali-Bashi, Maryam Peymani, Motahare-Sadat Hashemi, Masoud Etemadifar, Mohammad Hossein Nasr-Esfahani, Kamran Ghaedi

There is an error in affiliation 2 for author Maryam Peymani. The correct affiliation 2 is: Department of Biology, Faculty of Basic Sciences, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran.

Reference

Fazeli S, Motovali-Bashi M, Peymani M, Hashemi M- S, Etemadifar M, Nasr-Esfahani MH, et al. (2020)
 A compound downregulation of SRRM2 and miR-27a-3p with upregulation of miR-27b-3p in PBMCs of
 Parkinson's patients is associated with the early stage onset of disease. PLoS ONE 15(11): e0240855.
 https://doi.org/10.1371/journal.pone.0240855 PMID: 33171483



GOPEN ACCESS

Citation: Fazeli S, Motovali-Bashi M, Peymani M, Hashemi M-S, Etemadifar M, Hossein Nasr-Esfahani M, et al. (2020) Correction: A compound downregulation of *SRRM2* and miR-27a-3p with upregulation of miR-27b-3p in PBMCs of Parkinson's patients is associated with the early stage onset of disease. PLoS ONE 15(12): e0244776. https://doi.org/10.1371/journal.pone.0244776

Published: December 30, 2020

Copyright: © 2020 Fazeli et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.